Neuroscience Letters, 217 (1996) 212–214
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Abe, K., see Sakurai, M., 113
Agapito, M.T., see Pablos, M.I., 161
Agnati, L.F., see Cintra, A., 61
Aistrup, G.L., see Nagata, K., 189
Akaike, A., see Fukuda, K., 13
Aksenov, M.Y., Aksenova, M.V., Carney, J.M., Butterfield, D.A., α1-Antichymotrypsin interaction with Aβ(1–42) does not inhibit fibril formation but attenuates the peptide toxicity, 117
Aksenova, M.V., see Aksenov, M.Y., 117
Amant, C., see Amouyel, P., 203
Amenta, F., see Sabbatini, M., 85
Amouyel, P., Richard, F., Cottel, D., Amant, C., Codron, V., Helbecque, N., The deletion allele of the angiotensin I converting enzyme gene as a genetic susceptibility factor for cognitive impairment, 203
Andbjer, B., see Cintra, A., 61
Aoki, M., see Sakurai, M., 113
Araki, T., see Taniwaki, Y., 29
Arranz, M.J., Collier, D.A., Munro, J., Sham, P., Kirov, G., Sodhi, M., Roberts, G., Price, J., Kerwin, R.W., Analysis of a structural polymorphism in the 5-HT_{2A} receptor and clinical response to clozapine, 177

Baek, S.Y., see Park, H.T., 17
Bahr, B.A., see Hall, R.A., 179
Bajaj, N.P.S., see Guidato, S., 157
Baker, I., see Masserano, J.M., 73
Bannon, M.J., see Moody, C.A., 55
Baran, L., see Przegaliński, E., 145
Barthélémy, P., Sabeur, G., Jammes, Y., Reflex relationships between the cervical esophagus and the respiratory system in cats, 81
Blomqvist, A., see Hamlin, L., 149
Bourdel, E., see Van Kemmel, F.-M., 58
Brecha, N.C., see De Giorgio, R., 77
Butterfield, D.A., see Aksenov, M.Y., 117

Camón, L., see Giménez-Llort, L., 1
Carney, J.M., see Aksenov, M.Y., 117
Choi, R.C.Y., Leung, P.W.Y., Dong, T.T.X., Wan, D.C.C., Tsim, K.W.K., Calcitonin gene-related peptide increases the expression of acetylcholinesterase in cultured chick myotubes, 165
Cintra, A., Andbjer, B., Finnman, U.-B., Hagman, M., Agnati, L.F., Höglund, G., Fuxe, K., Subacute toluene exposure increases DA dysfunction in the 6-OH dopamine lesioned nigrostriatal dopaminergic system of the rat, 61
Codron, V., see Amouyel, P., 203
Collier, D.A., see Arranz, M.J., 177
Costentin, J., see Van Kemmel, F.-M., 58
Cottel, D., see Amouyel, P., 203

De Giorgio, R., Su, D., Peter, D., Edwards, R.H., Brecha, N.C., Sternini, C., Vesicular monoamine transporter 2 expression in enteric neurons and enterochromaffin-like cells of the rat, 77
de Vera, N., see Giménez-Llort, L., 1
Deppe, M., see Knecht, S., 66
Diehl, B., see Knecht, S., 66
Dong, T.T.X., see Choi, R.C.Y., 165
Doutheil, J., see Paschen, W., 41
Dubuc, I., see Van Kemmel, F.-M., 58

Edwards, R.H., see De Giorgio, R., 77
Eisenhauer, P.B., see Wells, J.M., 121
Engelmann, M., see Liebsch, G., 101
Ericson, A.-C., see Hamlin, L., 149

Fehrentz, J.-A., see Van Kemmel, F.-M., 58
Fine, R.E., see Wells, J.M., 121
Finnman, U.-B., see Cintra, A., 61
Forsby, A., Walum, E., Polygodial induces inositol phosphate turnover in human neuroblastoma SH-SY5Y cells, 50
Fukuda, K., Kaneko, S., Yada, N., Kikuwaka, M., Akaike, A., Satoh, M., Cyclic AMP-dependent modulation of N- and Q-type Ca^{2+} channels expressed in *Xenopus* oocytes, 13
Fuxe, K., see Cintra, A., 61

Giménez-Llort, L., Martínez, E., Camón, L., de Vera, N., Concentration of putrescine in plasma, frontal cortex and hippocampus of rats after systemic administration of the convulsants *N*-methyl-D-aspartate, pentylentetrazol, picrotoxinine, lindane and 4-aminopyridine, 1
Gissel, C., see Paschen, W., 41
Goguen, D., see Guido, M.E., 9
Golan, H., Talpalar, A.E., Schleifstein-Attias, D., Grossman, Y., GABA metabolism controls inhibition efficacy in the mammalian CNS, 25
Granneman, J.G., see Moody, C.A., 55
Grossman, Y., see Golan, H., 25
Guidato, S., Bajaj, N.P.S., Miller, C.C.J., Cellular phosphorylation of neurofilament heavy-chain by cyclin-dependent kinase-5 masks the epitope for monoclonal antibody N52, 157
Guido, M.E., Goguen, D., Robertson, H.A., Rusak, B., Spontaneous and light-evoked expression of JunB-like protein in the hamster suprachiasmatic nucleus near subjective dawn, 9
Gutierrez-Baraja, R., see Pablos, M.I., 161

Hadjiconstantinou, M., Neff, N.H., Zhou, L.-W., Weiss, B., D_2 dopamine receptor antisense increases the activity and mRNA of tyrosine hydroxylase and aromatic L-amino acid decarboxylase in mouse brain, 105
Hagman, M., see Cintra, A., 61
Hall, R.A., Vodyanoy, V., Quan, A., Sinnarajah, S., Suppiramaniam, V., Kessler, M., Bahr, B.A., Effects of heparin on the properties of solubilized and reconstituted rat brain AMPA receptors, 179
Hamlin, L., Mackerlova, L., Blomqvist, A., Ericson, A.-C., AMPA-selective glutamate receptor subunits and their relation to glutamate- and GABA-like immunoreactive terminals in the nucleus submedius of the rat, 149
Helbecque, N., see Amouyel, P., 203
Henningsen, H., see Knecht, S., 66
Higuchi, T., see Okere, C.O., 197
Hirai, K., see Ohta, H., 37
Hirobe, M., see Kotake, Y., 69
Höglund, G., see Cintra, A., 61
Horiuchi, S., see Kimura, T., 209

Neuroscience Letters, 217 (1996) 212–214
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Abe, K., see Sakurai, M., 113
Agapito, M.T., see Pablos, M.I., 161
Agnati, L.F., see Cintra, A., 61
Aistrup, G.L., see Nagata, K., 189
Akaike, A., see Fukuda, K., 13
Aksenov, M.Y., Aksenova, M.V., Carney, J.M., Butterfield, D.A., α1-Antichymotrypsin interaction with Aβ(1–42) does not inhibit fibril formation but attenuates the peptide toxicity, 117
Aksenova, M.V., see Aksenov, M.Y., 117
Amant, C., see Amouyel, P., 203
Amenta, F., see Sabbatini, M., 85
Amouyel, P., Richard, F., Cottel, D., Amant, C., Codron, V., Helbecque, N., The deletion allele of the angiotensin I converting enzyme gene as a genetic susceptibility factor for cognitive impairment, 203
Andbjer, B., see Cintra, A., 61
Aoki, M., see Sakurai, M., 113
Araki, T., see Taniwaki, Y., 29
Arranz, M.J., Collier, D.A., Munro, J., Sham, P., Kirov, G., Sodhi, M., Roberts, G., Price, J., Kerwin, R.W., Analysis of a structural polymorphism in the 5-HT_{2A} receptor and clinical response to clozapine, 177

Baek, S.Y., see Park, H.T., 17
Bahr, B.A., see Hall, R.A., 179
Bajaj, N.P.S., see Guidato, S., 157
Baker, I., see Masserano, J.M., 73
Bannon, M.J., see Moody, C.A., 55
Baran, L., see Przegaliński, E., 145
Barthélémy, P., Sabeur, G., Jammes, Y., Reflex relationships between the cervical esophagus and the respiratory system in cats, 81
Blomqvist, A., see Hamlin, L., 149
Bourdel, E., see Van Kemmel, F.-M., 58
Brecha, N.C., see De Giorgio, R., 77
Butterfield, D.A., see Aksenov, M.Y., 117

Camón, L., see Giménez-Llort, L., 1
Carney, J.M., see Aksenov, M.Y., 117
Choi, R.C.Y., Leung, P.W.Y., Dong, T.T.X., Wan, D.C.C., Tsim, K.W.K., Calcitonin gene-related peptide increases the expression of acetylcholinesterase in cultured chick myotubes, 165
Cintra, A., Andbjer, B., Finnman, U.-B., Hagman, M., Agnati, L.F., Höglund, G., Fuxe, K., Subacute toluene exposure increases DA dysfunction in the 6-OH dopamine lesioned nigrostriatal dopaminergic system of the rat, 61
Codron, V., see Amouyel, P., 203
Collier, D.A., see Arranz, M.J., 177
Costentin, J., see Van Kemmel, F.-M., 58
Cottel, D., see Amouyel, P., 203

De Giorgio, R., Su, D., Peter, D., Edwards, R.H., Brecha, N.C., Sternini, C., Vesicular monoamine transporter 2 expression in enteric neurons and enterochromaffin-like cells of the rat, 77
de Vera, N., see Giménez-Llort, L., 1
Deppe, M., see Knecht, S., 66
Diehl, B., see Knecht, S., 66
Dong, T.T.X., see Choi, R.C.Y., 165
Doutheil, J., see Paschen, W., 41
Dubuc, I., see Van Kemmel, F.-M., 58

Edwards, R.H., see De Giorgio, R., 77
Eisenhauer, P.B., see Wells, J.M., 121
Engelmann, M., see Liebsch, G., 101
Ericson, A.-C., see Hamlin, L., 149

Fehrentz, J.-A., see Van Kemmel, F.-M., 58
Fine, R.E., see Wells, J.M., 121
Finnman, U.-B., see Cintra, A., 61
Forsby, A., Walum, E., Polygodial induces inositol phosphate turnover in human neuroblastoma SH-SY5Y cells, 50
Fukuda, K., Kaneko, S., Yada, N., Kikuwaka, M., Akaike, A., Satoh, M., Cyclic AMP-dependent modulation of N- and Q-type Ca^{2+} channels expressed in *Xenopus* oocytes, 13
Fuxe, K., see Cintra, A., 61

Giménez-Llort, L., Martínez, E., Camón, L., de Vera, N., Concentration of putrescine in plasma, frontal cortex and hippocampus of rats after systemic administration of the convulsants *N*-methyl-D-aspartate, pentylentetrazol, picrotoxinine, lindane and 4-aminopyridine, 1
Gissel, C., see Paschen, W., 41
Goguen, D., see Guido, M.E., 9
Golan, H., Talpalar, A.E., Schleifstein-Attias, D., Grossman, Y., GABA metabolism controls inhibition efficacy in the mammalian CNS, 25
Granneman, J.G., see Moody, C.A., 55
Grossman, Y., see Golan, H., 25
Guidato, S., Bajaj, N.P.S., Miller, C.C.J., Cellular phosphorylation of neurofilament heavy-chain by cyclin-dependent kinase-5 masks the epitope for monoclonal antibody N52, 157
Guido, M.E., Goguen, D., Robertson, H.A., Rusak, B., Spontaneous and light-evoked expression of JunB-like protein in the hamster suprachiasmatic nucleus near subjective dawn, 9
Gutierrez-Baraja, R., see Pablos, M.I., 161

Hadjiconstantinou, M., Neff, N.H., Zhou, L.-W., Weiss, B., D_2 dopamine receptor antisense increases the activity and mRNA of tyrosine hydroxylase and aromatic L-amino acid decarboxylase in mouse brain, 105
Hagman, M., see Cintra, A., 61
Hall, R.A., Vodyanoy, V., Quan, A., Sinnarajah, S., Suppiramaniam, V., Kessler, M., Bahr, B.A., Effects of heparin on the properties of solubilized and reconstituted rat brain AMPA receptors, 179
Hamlin, L., Mackerlova, L., Blomqvist, A., Ericson, A.-C., AMPA-selective glutamate receptor subunits and their relation to glutamate- and GABA-like immunoreactive terminals in the nucleus submedius of the rat, 149
Helbecque, N., see Amouyel, P., 203
Henningsen, H., see Knecht, S., 66
Higuchi, T., see Okere, C.O., 197
Hirai, K., see Ohta, H., 37
Hirobe, M., see Kotake, Y., 69
Höglund, G., see Cintra, A., 61
Horiuchi, S., see Kimura, T., 209

Iinuma, K., see Yokoyama, H., 194

Jammes, Y., see Barthélémy, P., 81
Justice Jr., J.B., see Welch, S.M., 184

Kaba, H., see Okere, C.O., 197
Kagitani, F., Kimura, A., Sato, A., Suzuki, A., The role of the spinal cord as a reflex center for the somatically induced reflex responses of splenic sympathetic and natural killer cell activity in anesthetized rats, 109
Kaneko, S., see Fukuda, K., 13
Kato, K., see Ohta, H., 37
Kato, M., see Taniwaki, Y., 29
Kaufmann, R., see Kinze, S., 45
Kerwin, R.W., see Arranz, M.J., 177
Kessler, M., see Hall, R.A., 179
Kikuwaka, M., see Fukuda, K., 13
Kim, B.S., see Park, H.T., 17
Kim, J.B., see Park, H.T., 17
Kim, J.J., see Park, H.T., 17
Kimura, A., see Kagitani, F., 109
Kimura, T., Horiuchi, S., N^{ε}-carboxymethyllysine and neurofibrillary tangles, 209
Kinze, S., Schöneberg, T., Meyer, R., Martin, H., Kaufmann, R., Pharmacological characterization of CCK_B receptors in human brain: no evidence for receptor heterogeneity, 45
Kirov, G., see Arranz, M.J., 177
Kishi, A., see Ohno, M., 21
Knecht, S., Henningsen, H., Deppe, M., Osinska, L., Diehl, B., Stodieck, S., Ringelstein, E.-B., Persistent unihemispheric perceptual impairments in humans following focal seizures, 66
Kobayashi, T., see Taniwaki, Y., 29
Kobayashi, Y., Matsumura, G., Central projections of primary afferent fibers from the rat trigeminal nerve labeled with isolectin B4-HRP, 89
Konishi, S., see Mitoma, H., 141
Korf, H.-W., see Stehle, J.H., 169
Korf, H.-W., see Weigle, C., 173
Kotake, Y., Yoshida, M., Ogawa, M., Tasaki, Y., Hirobe, M., Ohta, S., Chronic administration of 1-benzyl-1,2,3,4-tetrahydroisoquinoline, an endogenous amine in the brain, induces parkinsonism in a primate, 69
Kühn, R., see Stehle, J.H., 169

Landgraf, R., see Liebsch, G., 101
Leung, P.W.Y., see Choi, R.C.Y., 165
Liebsch, G., Wotjak, C.T., Landgraf, R., Engelmann, M., Septal vasopressin modulates anxiety-related behaviour in rats, 101
Luksch, H., Roth, G., Pretecto-tectal interactions: effects of lesioning and stimulating the pretectum on field potentials in the optic tectum of salamanders in vitro, 137
Luo, Y., Sunderland, T., Roth, G.S., Wolozin, B., Physiological levels of β-amyloid peptide promote PC12 cell proliferation, 125

Mackerlova, L., see Hamlin, L., 149
Marszalec, W., see Nagata, K., 189
Martin, H., see Kinze, S., 45
Martínez, E., see Giménez-Llort, L., 1
Martinez, J., see Van Kemmel, F.-M., 58
Masserano, J.M., Baker, I., Natsukari, N., Wyatt, R.J., Chronic cocaine administration increases tyrosine hydroxylase activity in the ventral tegmental area through glutaminergic- and dopaminergic D_2-receptor mechanisms, 73
Matsumura, G., see Kobayashi, Y., 89
McKenna, D.C., see Wells, J.M., 121
Meyer, R., see Kinze, S., 45
Miettinen, R., see Sorvari, H., 93
Miller, C.C.J., see Guidato, S., 157
Mitoma, H., Konishi, S., Long-lasting facilitation of inhibitory transmission by monoaminergic and cAMP-dependent mechanism in rat cerebellar GABAergic synapses, 141
Miyamoto, M., see Ohta, H., 37
Moody, C.A., Granneman, J.G., Bannon, M.J., Dopamine transporter binding in rat striatum and nucleus accumbens is unaltered following chronic changes in dopamine levels, 55
Munro, J., see Arranz, M.J., 177
Murphy Jr., G.M., Yang, L., Yesavage, J., Tinklenberg, J.R., Rate of cognitive decline in Alzheimer's disease is not affected by the alpha-1-antichymotrypsin A allele or the CYP2D6 B mutant, 200

Nagata, K., Aistrup, G.L., Huang, C.-S., Marszalec, W., Song, J.-H., Yeh, J.Z., Narahashi, T., Potent modulation of neuronal nicotinic acetylcholine receptor-channel by ethanol, 189
Narahashi, T., see Nagata, K., 189
Natsukari, N., see Masserano, J.M., 73
Neff, N.H., see Hadjiconstantinou, M., 105
Nishikawa, H., see Ohta, H., 37N

Ogawa, M., see Kotake, Y., 69
Oh, Y.-S., see Won, M.-H., 129
Ohno, M., Kishi, A., Watanabe, S., Effect of cholinergic activation by physostigmine on working memory failure caused in rats by pharmacological manipulation of hippocampal glutamatergic and 5-HTergic neurotransmission, 21
Ohta, H., Nishikawa, H., Hirai, K., Kato, K., Miyamoto, M., Relationship of impaired brain glucose metabolism to learning deficit in the senescence-accelerated mouse, 37
Ohta, S., see Kotake, Y., 69
Okere, C.O., Kaba, H., Takahashi, S., Higuchi, T., Intrabulbar infusions of (±)*S*-nitroso-*N*-acetylpenicillamine and L-arginine induce functional plasticity in the accessory olfactory bulb of female mice, 197
Onodera, K., see Yokoyama, H., 194
Osinska, L., see Knecht, S., 66

Pablos, M.I., Agapito, M.T., Gutierrez-Baraja, R., Reiter, R.J., Recio, J.M., Effect of calcium on melatonin secretion in chick pineal gland I, 161
Park, H.T., Baek, S.Y., Kim, B.S., Kim, J.B., Kim, J.J., Developmental expression of 'RZRβ, a putative nuclear-melatonin receptor' mRNA in the suprachiasmatic nucleus of the rat, 17
Paschen, W., Doutheil, J., Uto, A., Gissel, C., Changes in endoplasmic reticulum Ca^{2+}-ATPase mRNA levels in transient cerebral ischemia of rat: a quantitative polymerase chain reaction study, 41
Perry, G., see Smith, M.A., 210
Perry, M.J.M., Alteration to the levels of galanin in uninjured contralateral rat saphenous nerves after unilateral saphenous nerve section, 206
Peter, D., see De Giorgio, R., 77
Pezeshki, G., Schöbitz, B., Pohl, T., Reul, J.M.H.M., Intracerebroventricular administration of missense oligodeoxynucleotide induces interleukin-6 mRNA expression in brain and spleen of rats, 97
Pfeffer, M., see Stehle, J.H., 169

Pitkänen, A., see Sorvari, H., 93
Pohl, T., see Pezeshki, G., 97
Price, J., see Arranz, M.J., 177
Przegaliński, E., Baran, L., Siwanowicz, J., The role of nitric oxide in chemically- and electrically-induced seizures in mice, 145

Quan, A., see Hall, R.A., 179

Recio, J.M., see Pablos, M.I., 161
Reiter, R.J., see Pablos, M.I., 161
Reul, J.M.H.M., see Pezeshki, G., 97
Richard, F., see Amouyel, P., 203
Ringelstein, E.-B., see Knecht, S., 66
Roberts, G., see Arranz, M.J., 177
Robertson, H.A., see Guido, M.E., 9
Rodman, H.R., see Sorenson, K.M., 5
Roth, G., see Luksch, H., 137
Roth, G.S., see Luo, Y., 125
Rusak, B., see Guido, M.E., 9

Sabbatini, M., Vega, J.A., Amenta, F., Peripheral nerve vascular changes in spontaneously hypertensive rats, 85
Sabeur, G., see Barthélémy, P., 81
Sadahiro, M., see Sakurai, M., 113
Sakurai, M., Aoki, M., Abe, K., Sadahiro, M., Tabayashi, K., Dissociation of HSP72 and HSC73 heat shock mRNA inductions after spinal cord ischemia in rabbit, 113
Sato, A., see Kagitani, F., 109
Sato, M., see Yokoyama, H., 194
Satoh, M., see Fukuda, K., 13
Schleifstein-Attias, D., see Golan, H., 25
Schöbitz, B., see Pezeshki, G., 97
Schöneberg, T., see Kinze, S., 45
Sham, P., see Arranz, M.J., 177
Shin, H.-C., see Won, M.-H., 129
Shinoda, Y., see Sugihara, I., 33
Sinnarajah, S., see Hall, R.A., 179
Siwanowicz, J., see Przegaliński, E., 145
Smith, M.A., Tabaton, M., Perry, G., Early contribution of oxidative glycation in Alzheimer disease, 210
Sodhi, M., see Arranz, M.J., 177
Soininen, H., see Sorvari, H., 93
Song, J.-H., see Nagata, K., 189
Sorenson, K.M., Rodman, H.R., The lateral geniculate nucleus does not project to area TE in infant or adult macaques, 5
Sorvari, H., Miettinen, R., Soininen, H., Pitkänen, A., Parvalbumin-immunoreactive neurons make inhibitory synapses on pyramidal cells in the human amygdala: a light and electron microscopic study, 93
Stehle, J.H., Pfeffer, M., Kühn, R., Korf, H.-W., Light-induced expression of transcription factor ICER (inducible cAMP early repressor) in rat suprachiasmatic nucleus is phase-restricted, 169
Sternini, C., see De Giorgio, R., 77
Stodieck, S., see Knecht, S., 66
Su, D., see De Giorgio, R., 77
Sugihara, I., Wu, H., Shinoda, Y., Morphology of axon collaterals of single climbing fibers in the deep cerebellar nuclei of the rat, 33
Sunderland, T., see Luo, Y., 125
Suppiramaniam, V., see Hall, R.A., 179
Suzuki, A., see Kagitani, F., 109

Tabaton, M., see Smith, M.A., 210
Tabayashi, K., see Sakurai, M., 113
Takahashi, H., see Wakabayashi, K., 133
Takahashi, S., see Okere, C.O., 197
Talpalar, A.E., see Golan, H., 25
Taniwaki, Y., Kato, M., Araki, T., Kobayashi, T., Microglial activation by epileptic activities through the propagation pathway of kainic acid-induced hippocampal seizures in the rat, 29
Tasaki, Y., see Kotake, Y., 69
Thompson, S.W.N., Wall, P.D., The effect of GABA and 5-HT receptor antagonists on rat dorsal root potentials, 153
Tinklenberg, J.R., see Murphy Jr., G.M., 200
Tsim, K.W.K., see Choi, R.C.Y., 165

Ullman, M.D., see Wells, J.M., 121
Uto, A., see Paschen, W., 41

Van Kemmel, F.-M., Dubuc, I., Bourdel, E., Fehrentz, J.-A., Martinez, J., Costentin, J., A C-terminal cyclic 8–13 neurotensin fragment analog appears less exposed to neprilysin when it crosses the blood–brain barrier than the cerebrospinal fluidbrain barrier in mice, 58
Vega, J.A., see Sabbatini, M., 85
Ventura, R.F., see Wells, J.M., 121
Vodyanoy, V., see Hall, R.A., 179

Wakabayashi, K., Takahashi, H., Gallyas-positive, tau-negative glial inclusions in Parkinson's disease midbrain, 133
Wall, P.D., see Thompson, S.W.N., 153
Walum, E., see Forsby, A., 50
Wan, D.C.C., see Choi, R.C.Y., 165
Watanabe, S., see Ohno, M., 21
Watanabe, T., see Yokoyama, H., 194
Weigle, C., Wicht, H., Korf, H.-W., A possible homologue of the suprachiasmatic nucleus in the hypothalamus of lampreys (*Lampetra fluviatilis* L.), 173
Weiss, B., see Hadjiconstantinou, M., 105
Welch, S.M., Justice Jr., J.B., Regulation of dopamine uptake in rat striatal tissue by NMDA receptors as measured using rotating disk electrode voltammetry, 184
Wells, J.M., Ventura, R.F., Eisenhauer, P.B., McKenna, D.C., Fine, R.E., Ullman, M.D., Transport of GM1 and GM1 inner ester across an in vitro model of the blood–brain barrier, 121
Wicht, H., see Weigle, C., 173
Wolozin, B., see Luo, Y., 125
Won, M.-H., Oh, Y.-S., Shin, H.-C., Localization of somatostatin-like immunoreactive neurons in the vestibular ganglion of the rabbit, 129
Wotjak, C.T., see Liebsch, G., 101
Wu, H., see Sugihara, I., 33
Wyatt, R.J., see Masserano, J.M., 73

Yada, N., see Fukuda, K., 13
Yang, L., see Murphy Jr., G.M., 200
Yeh, J.Z., see Nagata, K., 189
Yesavage, J., see Murphy Jr., G.M., 200
Yokoyama, H., Sato, M., Iinuma, K., Onodera, K., Watanabe, T., Centrally acting histamine H_1 antagonists promote the development of amygdala kindling in rats, 194
Yoshida, M., see Kotake, Y., 69
Zhou, L.-W., see Hadjiconstantinou, M., 105